



# Chlordiazepoxide

(Pronounced claw-dye-az-ee-pox-ide)

## Why has chlordiazepoxide been prescribed?

Chlordiazepoxide is known as an anxiolytic and also a hypnotic drug (sleeping tablet). These medications are prescribed to treat anxiety, insomnia and a variety of other conditions. It is also used as a treatment in alcohol dependence, to relieve symptoms of alcohol withdrawal.

## What are the benefits of taking chlordiazepoxide?

Chlordiazepoxide may be helpful in relieving anxiety, relaxing muscles and tension and in helping people who have difficulty getting to sleep.

## Are there any precautions with chlordiazepoxide?

Chlordiazepoxide is suitable for most people. As with all medicines, however, there are precautions. Your prescriber will usually check that it is safe to prescribe chlordiazepoxide, but let your prescriber know if any of the following apply to you, as extra care may be needed:

- a) If you have myasthenia gravis, sleep apnoea syndrome, severe lung disease, respiratory failure, kidney or liver trouble.
- b) If you are taking any other medication, such as cimetidine, sedative antihistamines, antibiotics such as rifampicin, medication for Parkinson's disease, medication for schizophrenia, medication for anxiety or insomnia, antidepressants, anticonvulsants or strong pain killers. Also, tell your prescriber if you buy any medicine 'over the counter' from a pharmacy or supermarket. Please also tell your prescriber if you take any alternative or complementary medicine such as Chinese herbal medicines.
- c) If you are pregnant, breast feeding, or wish to become pregnant.
- d) If you have lactose intolerance as the tablets contain lactose.

## What is the usual dose of chlordiazepoxide?

For anxiety the usual dose is 10mg three times daily. This may be increased up to 100mg in divided doses. For insomnia the dose may be between 5-15mg a day.

For alcohol withdrawal between 10-50mg four times a day up to a maximum of 250mg daily may be used depending on the severity of dependence.

Doses may be lower in older people.

## How should chlordiazepoxide be taken?

Chlordiazepoxide can be taken four times during the day for anxiety or add alcohol withdrawal, for insomnia, once a day, at night. However, you and your prescriber may decide that it is better for you to take the medicine in another way.

## What form does chlordiazepoxide come in?

Chlordiazepoxide is available only as capsules. The capsules are available as 5mg and 10mg.

## What should be done if a dose is missed?

**Anxiety:** If you forget a dose, take it as soon as you remember. If you miss a whole day's dose – just carry on as normal with the next day's dose.

**Insomnia:** Ideally chlordiazepoxide should not be taken regularly. It is recommended that it should only be taken every second or third night or when absolutely necessary.

**Alcohol withdrawal:** Take it as soon as you remember. If you are still experiencing excessive withdrawal symptoms after an hour of taking contact your prescriber.

## What will happen when chlordiazepoxide is first taken?

Chlordiazepoxide should start to work quite quickly, usually within 30 to 60 minutes. Many people do not experience any side effects. However some people may experience some side effects. We have listed potential side effects in the table below, whether or not they are likely to be short or long-term and what measures can be taken is also described. There are other possible side effects – we have listed only the most important ones. Ask your doctor, pharmacist or nurse if you are worried about anything else that you think might be a side effect. Further information on side effects is available in the official manufacturer's leaflet.

## Reporting side effects

The 'Yellow Card' scheme encourages patients to report any side effects that they feel may be caused by their medication. Ask your doctor, pharmacist, or nurse for the forms if you wish to report any side effects. Alternatively, telephone 0808 100 3352 or go to [www.mhra.gov.uk](http://www.mhra.gov.uk).

## Is chlordiazepoxide addictive?

Chlordiazepoxide may be addictive if it is taken continually for extended periods or at high doses. There may also be a risk that tolerance may develop. Tolerance means that higher doses are needed to achieve the same effect. If you stop taking the medicine suddenly, you may experience unpleasant physical feelings. Symptoms usually only occur after extended use and can include return of insomnia, anxiety, sweating, headache, palpitations and panic attacks. Reducing the dose of chlordiazepoxide slowly may make these symptoms less severe.

## What about alcohol?

It is recommended that people taking chlordiazepoxide should not drink alcohol. This is because both chlordiazepoxide and alcohol can cause drowsiness and impulsive behaviour. If the two are taken together, severe drowsiness or impulsiveness can result. This can lead to falls and accidents. As well as this, drinking alcohol may make anxiety and sleep problems worse. Never drink alcohol and drive. Discuss any concerns you have with your doctor, pharmacist or nurse.

## Is it OK to stop taking chlordiazepoxide when symptoms go away?

Yes. Chlordiazepoxide should be taken only for short periods - ideally for less than four weeks. Other methods of dealing with anxiety and insomnia may also be tried. These

include relaxation techniques and psychological therapies such as cognitive behavioural therapy. You should always discuss with your prescriber any plans you have to reduce or stop any of your prescribed medicines.

## Are there any alternatives to chlordiazepoxide?

Yes, there are alternatives available. Although anxiolytic and hypnotic drugs have broadly similar therapeutic effects they differ in the side effects they produce. There are other treatments for anxiety and insomnia available so most people can find one which suits them. Psychological and practical therapies can also help in some people with anxiety and insomnia, either as an alternative to medication or alongside it.

## Summary of side effects



### Drowsiness – Common (the morning after)

#### What can be done about it?

Do not drive if you are drowsy or dizzy the next day. Try to allow 7 to 8 hours for uninterrupted sleep after taking.

Other (less common) side effects	How common is it and what can be done about it?
Confusion	<b>Uncommon</b> Tell your prescriber if you experience confusion.
Muscle weakness	<b>Uncommon</b> Tell your prescriber if you experience muscle weakness.
Amnesia (memory loss)	<b>Uncommon</b> Tell your prescriber if taking chlordiazepoxide causes any memory loss.
Incoordination	<b>Uncommon</b> Tell your prescriber if you experience difficulty coordinating movements.
Dizziness	<b>Rare</b> This may wear off within a few days. Discuss with your prescriber if this persists.
Headache	<b>Rare</b> This should wear off within a few days. Paracetamol can be taken but discuss with your prescriber if this persists.
Dry mouth	<b>Rare</b> This should wear off within a few days. Sugar-free boiled sweets, chewing gum or

	eating citrus fruits may help. Discuss with the prescriber if this persists.
Disinhibition (loss of self control and inhibitions)	<b>Rare</b> Tell your prescriber if your behaviour changes in any way.
Depressed mood	<b>Rare</b> Tell your prescriber if taking chlordiazepoxide causes any changes in mood, perception, or thinking ability.
Irritability	<b>Rare</b> Tell your prescriber if taking chlordiazepoxide causes any changes in mood, perception, or thinking ability.
Aggression	<b>Rare</b> Tell your prescriber if taking chlordiazepoxide causes any changes in mood, perception, or thinking ability.
Nightmares	<b>Very rare</b> Tell your prescriber if taking chlordiazepoxide causes any changes in mood, perception, or thinking ability.
Hallucinations	<b>Very rare</b> Tell your prescriber if taking chlordiazepoxide causes any changes in mood, perception, or thinking ability.
Rash	<b>Very rare</b> Tell the prescriber straightaway if any rash develops.
Blood disorders	<b>Very rare</b> Any sore throat, fever, bruising or nose bleeds should be reported to your prescriber immediately
Jaundice	<b>Very rare</b> If you notice any yellowing of the skin or eyes contact your prescriber straightaway.

Very common = almost everyone affected

Common = many people affected

Uncommon = some people affected

Rare = few people affected

Very rare = very few or no one affected

## Disclaimer

This leaflet does not replace the official manufacturer's Patient Information Leaflet.

For more information, visit [www.slam.nhs.uk/meds](http://www.slam.nhs.uk/meds) or speak to a health care professional.

**Leaflet supplied by:** Medicines Information, Pharmacy Department, Maudsley Hospital, London SE5 8AZ | Telephone: 020 3228 2317

**Artwork by:** Oscar Millar ([www.millarstratton.co.uk](http://www.millarstratton.co.uk))

