



# Fluvoxamine

(Pronounced flew-vox-a-meen)

## Why has fluvoxamine been prescribed?

Fluvoxamine is an antidepressant known as a selective serotonin re-uptake inhibitor or SSRI. These medications are prescribed to treat depression and other mood disorders such as anxiety. Fluvoxamine is also used for obsessive compulsive disorder.

## What are the benefits of taking fluvoxamine?

Antidepressants are effective in reducing the severity of the symptoms of depression. They also shorten the time to recovery and prevent depression returning.

## Are there any precautions with fluvoxamine?

Fluvoxamine is suitable for most people. As with all medicines, however, there are precautions. Your prescriber will usually check that it is safe to prescribe fluvoxamine, but let your prescriber know if any of the following apply to you, as extra care may be needed:

- a) If you have epilepsy, diabetes, glaucoma, Parkinson's disease, a history of stomach ulcers or suffer from heart, liver or kidney trouble.
- b) If you are taking any other medication particularly anticoagulants such as warfarin, respiratory medicines such as theophylline, drugs for schizophrenia, bipolar disorder, depression and anti-inflammatory drugs or drugs for migraine. Also, tell your prescriber if you buy any medicine 'over the counter' from a pharmacy or supermarket. Please also tell your prescriber if you take any alternative or complementary medicine such as Chinese herbal medicines. St John's Wort should not be taken by patients prescribed fluvoxamine.
- c) If you are pregnant, breast feeding, or wish to become pregnant.

## What is the usual dose of fluvoxamine?

The starting dose is usually 50mg a day. The usual dose of fluvoxamine may be between 100mg and 300mg depending on the condition for which it has been prescribed. It may take some time to get to the dose that is most suitable.

## How should fluvoxamine be taken?

Fluvoxamine is usually taken once a day, in the evening (twice a day if the total dose is greater than 150mg). However, you and your prescriber may decide that it is better for you to take the medicine at another time.

## What form does fluvoxamine come in?

Fluvoxamine is available only as tablets. The tablets are available as 50mg and 100mg.

## What should be done if a dose is missed?

If you forget a dose, take it as soon as you remember. If you miss a whole day's dose – just carry on as normal with the next day's dose. If you miss more than a day's dose, speak to your prescriber.

## What will happen when fluvoxamine is first taken?

Fluvoxamine, like many medicines, does not work straight away. For example, it may take several days or even weeks for some symptoms to improve. To begin with, some people find that fluvoxamine may help them feel more alert and less slowed down. Later, (usually in two to three weeks) other symptoms should begin to improve. Many people do not

experience any side effects. However some people may experience side effects. We have listed potential side effects in the table below, whether or not they are likely to be short or long-term and what measures can be taken is also described. There are other possible side effects – we have listed only the most important ones. Ask your doctor, pharmacist or nurse if you are worried about anything else that you think might be a side effect. Further information on side effects is available in the official manufacturer’s leaflet.

## Reporting side effects

The ‘Yellow Card’ scheme encourages patients to report any side effects that they feel may be caused by their medication. Ask your doctor, pharmacist, or nurse for the forms if you wish to report any side effects. Alternatively, telephone 0808 100 3352 or go to [www.mhra.gov.uk](http://www.mhra.gov.uk).

## Is fluvoxamine addictive?

Fluvoxamine is not addictive, but if you stop taking the medicine suddenly, you may experience unpleasant physical feelings. Symptoms are fairly common and include nausea, insomnia, electric shock sensations, anxiety, dizziness and vivid dreams. Reducing the dose of fluvoxamine slowly may make these symptoms less severe.

## What about alcohol?

It is recommended that people taking fluvoxamine should not drink alcohol. This is because both antidepressants and alcohol can cause drowsiness. If the two are taken together, severe drowsiness can result. This can lead to falls and accidents. As well as this, drinking alcohol may make depression and anxiety worse. However, once people are used to taking medication, then *very small amounts* of alcohol may not be harmful. It pays to be very careful, because alcohol affects people in different ways, especially when they are taking medication. Never drink alcohol and drive. Discuss any concerns you may have with your doctor, pharmacist or nurse.

## Is it OK to stop taking fluvoxamine when symptoms go away?

No. If you stop taking fluvoxamine, your original symptoms are very likely to return. Most people need to remain on fluvoxamine for at least six months after they have fully recovered. You should always discuss with your prescriber any plans you have to reduce or stop any of your prescribed medicines.

## Are there any alternatives to fluvoxamine?

Yes, there are alternatives available. Although antidepressants have broadly similar therapeutic effects they differ in the side effects they produce. There are more than 20 antidepressants available so most people can find one which suits them. Psychological therapies can also help in some types of depression, either as an alternative to medication or alongside it.

## Summary of side effects



### Nausea - Common

#### What can be done about it?

This should wear off within a few days. Taking the medication with food may help. Discuss with your prescriber if this persists.



**Anxiety or restlessness – Common (feeling worried and agitated)**

**What can be done about it?**

This should wear off within a few days. Discuss with your prescriber if this persists.



**Insomnia or disturbed sleep - Common**

**What can be done about it?**

This should wear off within a few days. Fluvoxamine may be taken in the morning. Discuss with your prescriber if this persists.



**Diarrhoea – Common (loose stools)**

**What can be done about it?**

This should wear off within a few days. Discuss with your prescriber if this persists.



**Sexual dysfunction – Common (reduced libido (desire) and lack of orgasm in males and females. Impotence in males.)**

**What can be done about it?**

Any changes should be reported to your prescriber. It may be necessary to switch to another medication. Symptoms should go away when switched to another medication.

Other (less common) side effects	How common is it and what can be done about it?
Sweating	<b>Uncommon</b> Discuss this with your prescriber if it becomes troublesome.
Dry mouth	<b>Uncommon</b> This should wear off within a few days. Sugar-free boiled sweets, chewing gum or eating citrus fruits may help. Discuss with your prescriber if this persists.
Drowsiness	<b>Uncommon</b>

	This should wear off within a few days. Discuss with your prescriber if this persists. Taking the medication at a different time may help.
Headache	<b>Uncommon</b> This should wear off within a few days. Paracetamol can be taken but discuss with your prescriber if this persists.

Very common = almost everyone affected  
Common = many people affected  
Uncommon = some people affected

Rare = few people affected  
Very rare = very few or no one affected

## Disclaimer

This leaflet does not replace the official manufacturer's Patient Information Leaflet.

For more information, visit [www.slam.nhs.uk/meds](http://www.slam.nhs.uk/meds) or speak to a health care professional.

**Leaflet supplied by:** Medicines Information, Pharmacy Department, Maudsley Hospital, London SE5 8AZ | Telephone: 020 3228 2317

**Artwork by:** Oscar Millar ([www.millarstratton.co.uk](http://www.millarstratton.co.uk))